

Recommended Immunization Schedule for Persons Aged 0 Through 6 Years—United States • 2010

For those who fall behind or start late, see the catch-up schedule

Vaccine ▼	Age ►	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	19–23 months	2–3 years	4–6 years
Hepatitis B ¹		HepB	HepB			HepB						
Rotavirus ²			RV	RV	RV	RV ²						
Diphtheria, Tetanus, Pertussis ³			DTaP	DTaP	DTaP	<i>see footnote³</i>	DTaP					DTaP
<i>Haemophilus influenzae</i> type b ⁴			Hib	Hib	Hib ⁴	Hib						
Pneumococcal ⁵			PCV	PCV	PCV	PCV					PPSV	
Inactivated Poliovirus ⁶			IPV	IPV		IPV						IPV
Influenza ⁷						Influenza (Yearly)						
Measles, Mumps, Rubella ⁸							MMR			<i>see footnote⁸</i>		MMR
Varicella ⁹							Varicella			<i>see footnote⁹</i>		Varicella
Hepatitis A ¹⁰							HepA (2 doses)				HepA Series	
Meningococcal ¹¹												MCV

Range of recommended ages for all children except certain high-risk groups

Range of recommended ages for certain high-risk groups

This schedule includes recommendations in effect as of December 15, 2009. Any dose not administered at the recommended age should be administered at a subsequent visit, when indicated and feasible. The use of a combination vaccine generally is preferred over separate injections of its equivalent component vaccines. Considerations should include provider assessment, patient preference, and the potential for adverse events. Providers should consult the relevant Advisory

Committee on Immunization Practices statement for detailed recommendations: <http://www.cdc.gov/vaccines/pubs/acip-list.htm>. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS) at <http://www.vaers.hhs.gov> or by telephone, 800-822-7967.

1. Hepatitis B vaccine (HepB). (Minimum age: birth)

At birth:

- Administer monovalent HepB to all newborns before hospital discharge.
- If mother is hepatitis B surface antigen (HBsAg)-positive, administer HepB and 0.5 mL of hepatitis B immune globulin (HBIG) within 12 hours of birth.
- If mother's HBsAg status is unknown, administer HepB within 12 hours of birth. Determine mother's HBsAg status as soon as possible and, if HBsAg-positive, administer HBIG (no later than age 1 week).

After the birth dose:

- The HepB series should be completed with either monovalent HepB or a combination vaccine containing HepB. The second dose should be administered at age 1 or 2 months. Monovalent HepB vaccine should be used for doses administered before age 6 weeks. The final dose should be administered no earlier than age 24 weeks.
- Infants born to HBsAg-positive mothers should be tested for HBsAg and antibody to HBsAg 1 to 2 months after completion of at least 3 doses of the HepB series, at age 9 through 18 months (generally at the next well-child visit).
- Administration of 4 doses of HepB to infants is permissible when a combination vaccine containing HepB is administered after the birth dose. The fourth dose should be administered no earlier than age 24 weeks.

2. Rotavirus vaccine (RV). (Minimum age: 6 weeks)

- Administer the first dose at age 6 through 14 weeks (maximum age: 14 weeks 6 days). Vaccination should not be initiated for infants aged 15 weeks 0 days or older.
- The maximum age for the final dose in the series is 8 months 0 days
- If Rotarix is administered at ages 2 and 4 months, a dose at 6 months is not indicated.

3. Diphtheria and tetanus toxoids and acellular pertussis vaccine (DTaP). (Minimum age: 6 weeks)

- The fourth dose may be administered as early as age 12 months, provided at least 6 months have elapsed since the third dose.
- Administer the final dose in the series at age 4 through 6 years.

4. *Haemophilus influenzae* type b conjugate vaccine (Hib). (Minimum age: 6 weeks)

- If PRP-OMP (PedvaxHIB or Comvax [HepB-Hib]) is administered at ages 2 and 4 months, a dose at age 6 months is not indicated.
- TriHibit (DTaP/Hib) and Hiberix (PRP-T) should not be used for doses at ages 2, 4, or 6 months for the primary series but can be used as the final dose in children aged 12 months through 4 years.

5. Pneumococcal vaccine. (Minimum age: 6 weeks for pneumococcal conjugate vaccine [PCV]; 2 years for pneumococcal polysaccharide vaccine [PPSV])

- PCV is recommended for all children aged younger than 5 years. Administer 1 dose of PCV to all healthy children aged 24 through 59 months who are not completely vaccinated for their age.
- Administer PPSV 2 or more months after last dose of PCV to children aged 2 years or older with certain underlying medical conditions, including a cochlear implant. See *MMWR* 1997;46(No. RR-8).

6. Inactivated poliovirus vaccine (IPV) (Minimum age: 6 weeks)

- The final dose in the series should be administered on or after the fourth birthday and at least 6 months following the previous dose.
- If 4 doses are administered prior to age 4 years a fifth dose should be administered at age 4 through 6 years. See *MMWR* 2009;58(30):829–30.

7. Influenza vaccine (seasonal). (Minimum age: 6 months for trivalent inactivated influenza vaccine [TIV]; 2 years for live, attenuated influenza vaccine [LAIV])

- Administer annually to children aged 6 months through 18 years.
- For healthy children aged 2 through 6 years (i.e., those who do not have underlying medical conditions that predispose them to influenza complications), either LAIV or TIV may be used, except LAIV should not be given to children aged 2 through 4 years who have had wheezing in the past 12 months.
- Children receiving TIV should receive 0.25 mL if aged 6 through 35 months or 0.5 mL if aged 3 years or older.
- Administer 2 doses (separated by at least 4 weeks) to children aged younger than 9 years who are receiving influenza vaccine for the first time or who were vaccinated for the first time during the previous influenza season but only received 1 dose.
- For recommendations for use of influenza A (H1N1) 2009 monovalent vaccine see *MMWR* 2009;58(No. RR-10).

8. Measles, mumps, and rubella vaccine (MMR). (Minimum age: 12 months)

- Administer the second dose routinely at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 28 days have elapsed since the first dose.

9. Varicella vaccine. (Minimum age: 12 months)

- Administer the second dose routinely at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 3 months have elapsed since the first dose.
- For children aged 12 months through 12 years the minimum interval between doses is 3 months. However, if the second dose was administered at least 28 days after the first dose, it can be accepted as valid.

10. Hepatitis A vaccine (HepA). (Minimum age: 12 months)

- Administer to all children aged 1 year (i.e., aged 12 through 23 months). Administer 2 doses at least 6 months apart.
- Children not fully vaccinated by age 2 years can be vaccinated at subsequent visits
- HepA also is recommended for older children who live in areas where vaccination programs target older children, who are at increased risk for infection, or for whom immunity against hepatitis A is desired.

11. Meningococcal vaccine. (Minimum age: 2 years for meningococcal conjugate vaccine [MCV4] and for meningococcal polysaccharide vaccine [MPSV4])

- Administer MCV4 to children aged 2 through 10 years with persistent complement component deficiency, anatomic or functional asplenia, and certain other conditions placing them at high risk.
- Administer MCV4 to children previously vaccinated with MCV4 or MPSV4 after 3 years if first dose administered at age 2 through 6 years. See *MMWR* 2009;58:1042–3.

Recommended Immunization Schedule for Persons Aged 7 Through 18 Years—United States • 2010

For those who fall behind or start late, see the schedule below and the catch-up schedule

Vaccine ▼	Age ►	7–10 years	11–12 years	13–18 years	
Tetanus, Diphtheria, Pertussis ¹			Tdap	Tdap	Range of recommended ages for all children except certain high-risk groups
Human Papillomavirus ²	see footnote 2		HPV (3 doses)	HPV series	
Meningococcal ³		MCV	MCV	MCV	
Influenza ⁴			Influenza (Yearly)		Range of recommended ages for catch-up immunization
Pneumococcal ⁵			PPSV		
Hepatitis A ⁶			HepA Series		Range of recommended ages for certain high-risk groups
Hepatitis B ⁷			Hep B Series		
Inactivated Poliovirus ⁸			IPV Series		
Measles, Mumps, Rubella ⁹			MMR Series		
Varicella ¹⁰			Varicella Series		

This schedule includes recommendations in effect as of December 15, 2009. Any dose not administered at the recommended age should be administered at a subsequent visit, when indicated and feasible. The use of a combination vaccine generally is preferred over separate injections of its equivalent component vaccines. Considerations should include provider assessment, patient preference, and the potential for adverse events. Providers should consult the relevant Advisory

Committee on Immunization Practices statement for detailed recommendations: <http://www.cdc.gov/vaccines/pubs/acip-list.htm>. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS) at <http://www.vaers.hhs.gov> or by telephone, 800-822-7967.

1. Tetanus and diphtheria toxoids and acellular pertussis vaccine (Tdap). (Minimum age: 10 years for Boostrix and 11 years for Adacel)

- Administer at age 11 or 12 years for those who have completed the recommended childhood DTP/DTaP vaccination series and have not received a tetanus and diphtheria toxoid (Td) booster dose.
- Persons aged 13 through 18 years who have not received Tdap should receive a dose.
- A 5-year interval from the last Td dose is encouraged when Tdap is used as a booster dose; however, a shorter interval may be used if pertussis immunity is needed.

2. Human papillomavirus vaccine (HPV). (Minimum age: 9 years)

- Two HPV vaccines are licensed: a quadrivalent vaccine (HPV4) for the prevention of cervical, vaginal and vulvar cancers (in females) and genital warts (in females and males), and a bivalent vaccine (HPV2) for the prevention of cervical cancers in females.
- HPV vaccines are most effective for both males and females when given before exposure to HPV through sexual contact.
- HPV4 or HPV2 is recommended for the prevention of cervical precancers and cancers in females.
- HPV4 is recommended for the prevention of cervical, vaginal and vulvar precancers and cancers and genital warts in females.
- Administer the first dose to females at age 11 or 12 years.
- Administer the second dose 1 to 2 months after the first dose and the third dose 6 months after the first dose (at least 24 weeks after the first dose).
- Administer the series to females at age 13 through 18 years if not previously vaccinated.
- HPV4 may be administered in a 3-dose series to males aged 9 through 18 years to reduce their likelihood of acquiring genital warts.

3. Meningococcal conjugate vaccine (MCV4).

- Administer at age 11 or 12 years, or at age 13 through 18 years if not previously vaccinated.
- Administer to previously unvaccinated college freshmen living in a dormitory.
- Administer MCV4 to children aged 2 through 10 years with persistent complement component deficiency, anatomic or functional asplenia, or certain other conditions placing them at high risk.
- Administer to children previously vaccinated with MCV4 or MPSV4 who remain at increased risk after 3 years (if first dose administered at age 2 through 6 years) or after 5 years (if first dose administered at age 7 years or older). Persons whose only risk factor is living in on-campus housing are not recommended to receive an additional dose. See *MMWR* 2009;58:1042–3.

4. Influenza vaccine (seasonal).

- Administer annually to children aged 6 months through 18 years.
- For healthy nonpregnant persons aged 7 through 18 years (i.e., those who do not have underlying medical conditions that predispose them to influenza complications), either LAIV or TIV may be used.
- Administer 2 doses (separated by at least 4 weeks) to children aged younger than 9 years who are receiving influenza vaccine for the first time or who were vaccinated for the first time during the previous influenza season but only received 1 dose.
- For recommendations for use of influenza A (H1N1) 2009 monovalent vaccine. See *MMWR* 2009;58(No. RR-10).

5. Pneumococcal polysaccharide vaccine (PPSV).

- Administer to children with certain underlying medical conditions, including a cochlear implant. A single revaccination should be administered after 5 years to children with functional or anatomic asplenia or an immunocompromising condition. See *MMWR* 1997;46(No. RR-8).

6. Hepatitis A vaccine (HepA).

- Administer 2 doses at least 6 months apart.
- HepA is recommended for children aged older than 23 months who live in areas where vaccination programs target older children, who are at increased risk for infection, or for whom immunity against hepatitis A is desired.

7. Hepatitis B vaccine (HepB).

- Administer the 3-dose series to those not previously vaccinated.
- A 2-dose series (separated by at least 4 months) of adult formulation Recombivax HB is licensed for children aged 11 through 15 years.

8. Inactivated poliovirus vaccine (IPV).

- The final dose in the series should be administered on or after the fourth birthday and at least 6 months following the previous dose.
- If both OPV and IPV were administered as part of a series, a total of 4 doses should be administered, regardless of the child's current age.

9. Measles, mumps, and rubella vaccine (MMR).

- If not previously vaccinated, administer 2 doses or the second dose for those who have received only 1 dose, with at least 28 days between doses.

10. Varicella vaccine.

- For persons aged 7 through 18 years without evidence of immunity (see *MMWR* 2007;56[No. RR-4]), administer 2 doses if not previously vaccinated or the second dose if only 1 dose has been administered.
- For persons aged 7 through 12 years, the minimum interval between doses is 3 months. However, if the second dose was administered at least 28 days after the first dose, it can be accepted as valid.
- For persons aged 13 years and older, the minimum interval between doses is 28 days.

Recommended Adult Immunization Schedule


UNITED STATES - 2010


Note: These recommendations *must* be read with the footnotes that follow containing number of doses, intervals between doses, and other important information.

Figure 1. Recommended adult immunization schedule, by vaccine and age group

VACCINE ▼	AGE GROUP ▶	19–26 years	27–49 years	50–59 years	60–64 years	≥65 years
Tetanus, diphtheria, pertussis (Td/Tdap) ^{1,*}		Substitute 1-time dose of Tdap for Td booster; then boost with Td every 10 yrs				Td booster every 10 yrs
Human papillomavirus (HPV) ^{2,*}		3 doses (females)				
Varicella ^{3,*}		2 doses				
Zoster ⁴					1 dose	
Measles, mumps, rubella (MMR) ^{5,*}		1 or 2 doses		1 dose		
Influenza ^{6,*}		1 dose annually				
Pneumococcal (polysaccharide) ^{7,8}		1 or 2 doses				1 dose
Hepatitis A ^{9,*}		2 doses				
Hepatitis B ^{10,*}		3 doses				
Meningococcal ^{11,*}		1 or more doses				

*Covered by the Vaccine Injury Compensation Program.

 For all persons in this category who meet the age requirements and who lack evidence of immunity (e.g., lack documentation of vaccination or have no evidence of prior infection)

 Recommended if some other risk factor is present (e.g., on the basis of medical, occupational, lifestyle, or other indications)

 No recommendation

Report all clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System (VAERS). Reporting forms and instructions on filing a VAERS report are available at www.vaers.hhs.gov or by telephone, 800-822-7967.

Information on how to file a Vaccine Injury Compensation Program claim is available at www.hrsa.gov/vaccinecompensation or by telephone, 800-338-2382. To file a claim for vaccine injury, contact the U.S. Court of Federal Claims, 717 Madison Place, N.W., Washington, D.C. 20005; telephone, 202-357-6400.

Additional information about the vaccines in this schedule, extent of available data, and contraindications for vaccination is also available at www.cdc.gov/vaccines or from the CDC-INFO Contact Center at 800-CDC-INFO (800-232-4636) in English and Spanish, 24 hours a day, 7 days a week.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

Figure 2. Vaccines that might be indicated for adults based on medical and other indications

INDICATION ▶ VACCINE ▼	Pregnancy	Immuno-compromising conditions (excluding human immunodeficiency virus (HIV)) ^{3-5,13}	HIV infection ^{8-5,12,13}		Diabetes, heart disease, chronic lung disease, chronic alcoholism	Asplenia ¹² (including elective splenectomy and persistent complement deficiencies)	Chronic liver disease	Kidney failure, end-stage renal disease, receipt of hemodialysis	Health-care personnel		
			CD4+ T lymphocyte count <200 cells/μL	>200 cells/μL							
Tetanus, diphtheria, pertussis (Td/Tdap) ^{1,*}	Td	Substitute 1-time dose of Tdap for Td booster; then boost with Td every 10 yrs									
Human papillomavirus (HPV) ^{2,*}		3 doses for females through age 26 yrs									
Varicella ^{3,*}	Contraindicated				2 doses						
Zoster ⁴	Contraindicated				1 dose						
Measles, mumps, rubella (MMR) ^{5,*}	Contraindicated				1 or 2 doses						
Influenza ^{6,*}					1 dose TIV annually						
Pneumococcal (polysaccharide) ^{7,8}					1 or 2 doses						
Hepatitis A ^{9,*}					2 doses						
Hepatitis B ^{10,*}					3 doses						
Meningococcal ^{11,*}					1 or more doses				1 dose TIV or LAIV annually		

*Covered by the Vaccine Injury Compensation Program.

For all persons in this category who meet the age requirements and who lack evidence of immunity (e.g., lack documentation of vaccination or have no evidence of prior infection)
 Recommended if some other risk factor is present (e.g., on the basis of medical, occupational, lifestyle, or other indications)
 No recommendation

These schedules indicate the recommended age groups and medical indications for which administration of currently licensed vaccines is commonly indicated for adults ages 19 years and older, as of January 1, 2010. Licensed combination vaccines may be used whenever any components of the combination are indicated and when the vaccine's other components are not contraindicated. For detailed recommendations on all vaccines, including those used primarily for travelers or that are issued during the year, consult the manufacturers' package inserts and the complete statements from the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/pubs/acip-list.htm).

The recommendations in this schedule were approved by the Centers for Disease Control and Prevention's (CDC) Advisory Committee on Immunization Practices (ACIP), the American Academy of Family Physicians (AAFP), the American College of Obstetricians and Gynecologists (ACOG), and the American College of Physicians (ACP).



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



Footnotes

Recommended Adult Immunization Schedule—UNITED STATES - 2010

For complete statements by the Advisory Committee on Immunization Practices (ACIP), visit www.cdc.gov/vaccines/pubs/ACIP-list.htm.

1. Tetanus, diphtheria, and acellular pertussis (Td/Tdap) vaccination

Tdap should replace a single dose of Td for adults aged 19 through 64 years who have not received a dose of Tdap previously.

Adults with uncertain or incomplete history of primary vaccination series with tetanus and diphtheria toxoid-containing vaccines should begin or complete a primary vaccination series. A primary series for adults is 3 doses of tetanus and diphtheria toxoid-containing vaccines; administer the first 2 doses at least 4 weeks apart and the third dose 6–12 months after the second; Tdap can substitute for any one of the doses of Td in the 3-dose primary series. The booster dose of tetanus and diphtheria toxoid-containing vaccine should be administered to adults who have completed a primary series and if the last vaccination was received ≥ 10 years previously. Tdap or Td vaccine may be used, as indicated.

If a woman is pregnant and received the last Td vaccination ≥ 10 years previously, administer Td during the second or third trimester. If the woman received the last Td vaccination < 10 years previously, administer Tdap during the immediate postpartum period. A dose of Tdap is recommended for postpartum women, close contacts of infants aged < 12 months, and all health-care personnel with direct patient contact if they have not previously received Tdap. An interval as short as 2 years from the last Td is suggested; shorter intervals can be used. Td may be deferred during pregnancy and Tdap substituted in the immediate postpartum period, or Tdap can be administered instead of Td to a pregnant woman.

Consult the ACIP statement for recommendations for giving Td as prophylaxis in wound management.

2. Human papillomavirus (HPV) vaccination

HPV vaccination is recommended at age 11 or 12 years with catch-up vaccination at ages 13 through 26 years.

Ideally, vaccine should be administered before potential exposure to HPV through sexual activity; however, females who are sexually active should still be vaccinated consistent with age-based recommendations. Sexually active females who have not been infected with any of the four HPV vaccine types (types 6, 11, 16, 18 all of which HPV4 prevents) or any of the two HPV vaccine types (types 16 and 18 both of which HPV2 prevents) receive the full benefit of the vaccination. Vaccination is less beneficial for females who have already been infected with one or more of the HPV vaccine types. HPV4 or HPV2 can be administered to persons with a history of genital warts, abnormal Papanicolaou test, or positive HPV DNA test, because these conditions are not evidence of prior infection with all vaccine HPV types.

HPV4 may be administered to males aged 9 through 26 years to reduce their likelihood of acquiring genital warts. HPV4 would be most effective when administered before exposure to HPV through sexual contact.

A complete series for either HPV4 or HPV2 consists of 3 doses. The second dose should be administered 1–2 months after the first dose; the third dose should be administered 6 months after the first dose.

Although HPV vaccination is not specifically recommended for persons with the medical indications described in Figure 2, “Vaccines that might be indicated for adults based on medical and other indications,” it may be administered to these persons because the HPV vaccine is not a live-virus vaccine. However, the immune response and vaccine efficacy might be less for persons with the medical indications described in Figure 2 than in persons who do not have the medical indications described or who are immunocompetent. Health-care personnel are not at increased risk because of occupational exposure, and should be vaccinated consistent with age-based recommendations.

3. Varicella vaccination

All adults without evidence of immunity to varicella should receive 2 doses of single-antigen varicella vaccine if not previously vaccinated or the second dose if they have received only 1 dose, unless they have a medical contraindication. Special consideration should be given to those who 1) have close contact with persons at high risk for severe disease (e.g., health-care personnel and family contacts of persons with immunocompromising conditions) or 2) are at high risk for exposure or transmission (e.g., teachers; child-care employees; residents and staff members of institutional settings, including correctional institutions; college students; military personnel; adolescents and adults living in households with children; nonpregnant women of childbearing age; and international travelers).

Evidence of immunity to varicella in adults includes any of the following: 1) documentation of 2 doses of varicella vaccine at least 4 weeks apart; 2) U.S.-born before 1980 (although for health-care personnel and pregnant women, birth before 1980 should not be considered evidence of immunity); 3) history of varicella based on diagnosis or verification of varicella by a health-care provider (for a patient reporting a history of or presenting with an atypical case, a mild case, or both, health-care providers should seek either an epidemiologic link with a typical varicella case or to a laboratory-confirmed case or evidence of laboratory confirmation, if it was performed at the time of acute disease); 4) history of herpes zoster based on diagnosis or verification of herpes zoster by a health-care provider; or 5) laboratory evidence of immunity or laboratory confirmation of disease.

Pregnant women should be assessed for evidence of varicella immunity. Women who do not have evidence of immunity should receive the first dose of varicella vaccine upon completion or termination of pregnancy and before discharge from the health-care facility. The second dose should be administered 4–8 weeks after the first dose.

4. Herpes zoster vaccination

A single dose of zoster vaccine is recommended for adults aged ≥ 60 years regardless of whether they report a prior episode of herpes zoster. Persons with chronic medical conditions may be vaccinated unless their condition constitutes a contraindication.

5. Measles, mumps, rubella (MMR) vaccination

Adults born before 1957 generally are considered immune to measles and mumps.

Measles component: Adults born during or after 1957 should receive 1 or more doses of MMR vaccine unless they have 1) a medical contraindication; 2) documentation of vaccination with 1 or more doses of MMR vaccine; 3) laboratory evidence of immunity; or 4) documentation of physician-diagnosed measles.

A second dose of MMR vaccine, administered 4 weeks after the first dose, is recommended for adults who 1) have been recently exposed to measles or are in an outbreak setting; 2) have been vaccinated previously with killed measles vaccine; 3) have been vaccinated with an unknown type of measles vaccine during 1963–1967; 4) are students in postsecondary educational institutions; 5) work in a health-care facility; or 6) plan to travel internationally.

Mumps component: Adults born during or after 1957 should receive 1 dose of MMR vaccine unless they have 1) a medical contraindication; 2) documentation of vaccination with 1 or more doses of MMR vaccine; 3) laboratory evidence of immunity; or 4) documentation of physician-diagnosed mumps.

A second dose of MMR vaccine, administered 4 weeks after the first dose, is recommended for adults who 1) live in a community experiencing a mumps outbreak and are in an affected age group; 2) are students in postsecondary educational institutions; 3) work in a health-care facility; or 4) plan to travel internationally.

Rubella component: 1 dose of MMR vaccine is recommended for women who do not have documentation of rubella vaccination, or who lack laboratory evidence of immunity. For women of childbearing age, regardless of birth year, rubella immunity should be determined and women should be counseled regarding congenital rubella syndrome. Women who do not have evidence of immunity should receive MMR vaccine upon completion or termination of pregnancy and before discharge from the health-care facility.

Health-care personnel born before 1957: For unvaccinated health-care personnel born before 1957 who lack laboratory evidence of measles, mumps, and/or rubella immunity or laboratory confirmation of disease, health-care facilities should consider vaccinating personnel with 2 doses of MMR vaccine at the appropriate interval (for measles and mumps) and 1 dose of MMR vaccine (for rubella), respectively.

During outbreaks, health-care facilities should recommend that unvaccinated health-care personnel born before 1957, who lack laboratory evidence of measles, mumps, and/or rubella immunity or laboratory confirmation of disease, receive 2 doses of MMR vaccine during an outbreak of measles or mumps, and 1 dose during an outbreak of rubella.

Complete information about evidence of immunity is available at www.cdc.gov/vaccines/recs/provisional/default.htm.

6. Seasonal Influenza vaccination

Vaccinate all persons aged ≥ 50 years and any younger persons who would like to decrease their risk of getting influenza. Vaccinate persons aged 19 through 49 years with any of the following indications.

Medical: Chronic disorders of the cardiovascular or pulmonary systems, including asthma; chronic metabolic diseases, including diabetes mellitus; renal or hepatic dysfunction, hemoglobinopathies, or immunocompromising conditions (including immunocompromising conditions caused by medications or HIV); cognitive, neurologic or neuromuscular disorders; and pregnancy during the influenza season. No data exist on the risk for severe or complicated influenza disease among persons with asplenia; however, influenza is a risk factor for secondary bacterial infections that can cause severe disease among persons with asplenia.

Occupational: All health-care personnel, including those employed by long-term care and assisted-living facilities, and caregivers of children aged < 5 years.

Other: Residents of nursing homes and other long-term care and assisted-living facilities; persons likely to transmit influenza to persons at high risk (e.g., in-home household contacts and caregivers of children aged < 5 years, persons aged ≥ 50 years, and persons of all ages with high-risk conditions).

Healthy, nonpregnant adults aged < 50 years without high-risk medical conditions who are not contacts of severely immunocompromised persons in special-care units may receive either intranasally administered live, attenuated influenza vaccine (FluMist) or inactivated vaccine. Other persons should receive the inactivated vaccine.

7. Pneumococcal polysaccharide (PPSV) vaccination

Vaccinate all persons with the following indications.

Medical: Chronic lung disease (including asthma); chronic cardiovascular diseases; diabetes mellitus; chronic liver diseases, cirrhosis; chronic alcoholism; functional or anatomic asplenia (e.g., sickle cell disease or splenectomy [if elective splenectomy is planned, vaccinate at least 2 weeks before surgery]); immunocompromising conditions including chronic renal failure or nephrotic syndrome; and cochlear implants and cerebrospinal fluid leaks. Vaccinate as close to HIV diagnosis as possible.

Other: Residents of nursing homes or long-term care facilities and persons who smoke cigarettes. Routine use of PPSV is not recommended for American Indians/Alaska Natives or persons aged < 65 years unless they have underlying medical conditions that are PPSV indications. However, public health authorities may consider recommending PPSV for American Indians/Alaska Natives and persons aged 50 through 64 years who are living in areas where the risk for invasive pneumococcal disease is increased.

8. Revaccination with PPSV

One-time revaccination after 5 years is recommended for persons with chronic renal failure or nephrotic syndrome; functional or anatomic asplenia (e.g., sickle cell disease or splenectomy); and for persons with immunocompromising conditions. For persons aged ≥ 65 years, one-time revaccination is recommended if they were vaccinated ≥ 5 years previously and were younger than aged < 65 years at the time of primary vaccination.

9. Hepatitis A vaccination

Vaccinate persons with any of the following indications and any person seeking protection from hepatitis A virus (HAV) infection.

Behavioral: Men who have sex with men and persons who use injection drugs.

Occupational: Persons working with HAV-infected primates or with HAV in a research laboratory setting.

Medical: Persons with chronic liver disease and persons who receive clotting factor concentrates.

Other: Persons traveling to or working in countries that have high or intermediate endemicity of hepatitis A (a list of countries is available at www.cdc.gov/travel/content/diseases.aspx).

Unvaccinated persons who anticipate close personal contact (e.g., household contact or regular babysitting) with an international adoptee from a country of high or intermediate endemicity during the first 60 days after arrival of the adoptee in the United States should consider vaccination. The first dose of the 2-dose hepatitis A vaccine series should be administered as soon as adoption is planned, ideally ≥ 2 weeks before the arrival of the adoptee.

Single-antigen vaccine formulations should be administered in a 2-dose schedule at either 0 and 6–12 months (Havrix), or 0 and 6–18 months (Vaqta). If the combined hepatitis A and hepatitis B vaccine (Twinrix) is used, administer 3 doses at 0, 1, and 6 months; alternatively, a 4-dose schedule, administered on days 0, 7, and 21–30 followed by a booster dose at month 12 may be used.

10. Hepatitis B vaccination

Vaccinate persons with any of the following indications and any person seeking protection from hepatitis B virus (HBV) infection.

Behavioral: Sexually active persons who are not in a long-term, mutually monogamous relationship (e.g., persons with more than one sex partner during the previous 6 months); persons seeking evaluation or treatment for a sexually transmitted disease (STD); current or recent injection-drug users; and men who have sex with men.

Occupational: Health-care personnel and public-safety workers who are exposed to blood or other potentially infectious body fluids.

Medical: Persons with end-stage renal disease, including patients receiving hemodialysis; persons with HIV infection; and persons with chronic liver disease.

Other: Household contacts and sex partners of persons with chronic HBV infection; clients and staff members of institutions for persons with developmental disabilities; and international travelers to countries with high or intermediate prevalence of chronic HBV infection (a list of countries is available at www.cdc.gov/travel/content/diseases.aspx).

Hepatitis B vaccination is recommended for all adults in the following settings: STD treatment facilities; HIV testing and treatment facilities; facilities providing drug-abuse treatment and prevention services; health-care settings targeting services to injection-drug users or men who have sex with men; correctional facilities; end-stage renal disease programs and facilities for chronic hemodialysis patients; and institutions and nonresidential daycare facilities for persons with developmental disabilities.

Administer or complete a 3-dose series of HepB to those persons not previously vaccinated. The second dose should be administered 1 month after the first dose; the third dose should be administered at least 2 months after the second dose (and at least 4 months after the first dose). If the combined hepatitis A and hepatitis B vaccine (Twinrix) is used, administer 3 doses at 0, 1, and 6 months; alternatively, a 4-dose schedule, administered on days 0, 7, and 21–30 followed by a booster dose at month 12 may be used.

Adult patients receiving hemodialysis or with other immunocompromising conditions should receive 1 dose of 40 $\mu\text{g}/\text{mL}$ (Recombivax HB) administered on a 3-dose schedule or 2 doses of 20 $\mu\text{g}/\text{mL}$ (Engerix-B) administered simultaneously on a 4-dose schedule at 0, 1, 2 and 6 months.

11. Meningococcal vaccination

Meningococcal vaccine should be administered to persons with the following indications.

Medical: Adults with anatomic or functional asplenia, or persistent complement component deficiencies.

Other: First-year college students living in dormitories; microbiologists routinely exposed to isolates of *Neisseria meningitidis*; military recruits; and persons who travel to or live in countries in which meningococcal disease is hyperendemic or epidemic (e.g., the “meningitis belt” of sub-Saharan Africa during the dry season [December through June]), particularly if their contact with local populations will be prolonged. Vaccination is required by the government of Saudi Arabia for all travelers to Mecca during the annual Hajj.

Meningococcal conjugate vaccine (MCV4) is preferred for adults with any of the preceding indications who are aged ≤ 55 years; meningococcal polysaccharide vaccine (MPSV4) is preferred for adults aged ≥ 56 years. Revaccination with MCV4 after 5 years is recommended for adults previously vaccinated with MCV4 or MPSV4 who remain at increased risk for infection (e.g., adults with anatomic or functional asplenia). Persons whose only risk factor is living in on-campus housing are not recommended to receive an additional dose.

12. Selected conditions for which *Haemophilus influenzae* type b (Hib) vaccine may be used

Hib vaccine generally is not recommended for persons aged ≥ 5 years. No efficacy data are available on which to base a recommendation concerning use of Hib vaccine for older children and adults. However, studies suggest good immunogenicity in patients who have sickle cell disease, leukemia, or HIV infection or who have had a splenectomy. Administering 1 dose of Hib vaccine to these high-risk persons who have not previously received Hib vaccine is not contraindicated.

13. Immunocompromising conditions

Inactivated vaccines generally are acceptable (e.g., pneumococcal, meningococcal, influenza [inactivated influenza vaccine]) and live vaccines generally are avoided in persons with immune deficiencies or immunocompromising conditions. Information on specific conditions is available at www.cdc.gov/vaccines/pubs/acip-list.htm.